

Neural correlates of visual and motor decision processes

Jeffrey D Schall* and Narcisse P Bichot

Recent research has clarified and revealed characteristics of perceptual and motor decision processes in the brain. A democracy of sensory neurons discriminate the properties of a stimulus, while competition contrasts the attributes of stimuli across the visual field to locate conspicuous stimuli. Saliency and significance are weighed to select an object on which to focus attention and action. Experimentally combining neural and mental chronometry has determined the contribution of perceptual and motor processes to the duration and variability of behavioral reaction time. Whereas perceptual processing occupies a relatively constant amount of time for a given stimulus condition, the processes of mapping particular stimuli onto the appropriate behavior and preparing the motor response provide flexibility but introduce delay and variability in reaction time.

Addresses

Vanderbilt Vision Research Center, Department of Psychology,
Wilson Hall, Vanderbilt University, Nashville, Tennessee 37240, USA
*e-mail: jeffrey.d.schall@vanderbilt.edu

Current Opinion in Neurobiology 1998, **8**:211–217

<http://biomednet.com/elecref/0959438800800211>

© Current Biology Ltd ISSN 0959-4388

Abbreviations

ERP	event-related potential
FEF	frontal eye field
MT	middle temporal visual area
V1	primary visual cortex

Introduction

“To succumb to the preponderance of one set of influences over another set”, as Ambrose Bierce [1] characterized decision, may be an accurate summary of recent evidence from behavioral neurophysiology experiments. Although complex decisions are difficult to investigate, the decision of where to direct gaze and grasp has proven particularly accessible experimentally. Visual selection and attention have been reviewed recently [2–5], so we will emphasize work, framed by recent studies in our laboratory, on neural processes associated with saccade target selection and gaze control in the frontal eye field (FEF).

Vision with scanning eye movements

The decisions made by the visuomotor system of primates must be considered in light of the natural behaviors produced in ecological settings. To identify an object in a scene, the eyes move so that the image of the object falls on the fovea. Saccadic eye movements are the rapid shifts of gaze that redirect the visual axis onto a new point in the image. Saccades tend to direct gaze to conspicuous features in the scene, but knowledge and goals also strongly influence the focus of gaze [6]. Although the

relationship between cognitive states and the patterns of scanning eye movements has proven difficult to elucidate [7], recent studies have shown systematic relationships between gaze behavior and information acquisition for visually guided behavior [8].

Visual search tasks, employed for many years to investigate visual attention, have been used recently to investigate saccade target selection [9,10]. The results of these studies have provided strong evidence for distinguishing the process responsible for selecting the target where gaze will shift from the process regulating when gaze will shift [11••]. Other recent studies have demonstrated clearly that the focus of attention coincides with the target of an intended saccade [12,13•] and that directing attention can influence the production of saccades [14••,15]. These results indicate that a common visual selection mechanism governs both covert and overt orienting. This conclusion is supported by recent brain imaging studies showing that common regions in human frontal and parietal cortex are activated in association with attention and saccade tasks [16,17]. At the same time, it is important to remember that the relation between attention and eye movements is not obligatory. In fact, covert orienting may be little more than a state of visual selection without activating motor circuitry to produce overt orienting.

Normal visual behavior is accomplished through a continuous cycle of fixation and visual analysis interrupted by saccades. During fixation of one point in the image at least three processes take place. One process is the visual analysis of the image centered on the fovea to ascertain its identity. The other two are the visual analysis of the image in the periphery to locate potential targets for the next saccade, and the production of the saccade. Several key questions provide a framework for investigating the dynamics of visually guided behavior [18]. Namely, how much time does perceptual processing and response preparation take? Can these two processes overlap? How late can perceptual processing influence response preparation?

Neural correlates of saccade target selection in the FEF

Evolution of target selection signals

To investigate how the brain selects the target for a visually guided saccade, neural activity has been recorded in the FEF of monkeys making saccades to the oddball target in a popout visual search task [19,20••,21••,22•] or scanning complex images [23]. The FEF is an interesting area in which to investigate sensory decision processes because it is reciprocally connected with many extrastriate visual areas associated with both the dorsal and ventral visual streams [24], and also because it projects directly to

brainstem oculomotor structures [25,26]. Thus, the FEF is a nodal area, positioned to sample and possibly influence the outcome of visual processing and to play a central role in the production of eye movements [27,28•].

We found that the initial response of most visual neurons in the FEF did not discriminate whether the target or only distractors of the search array appeared in their receptive field (Figure 1a) [19,21••]. This observation should not be surprising because earlier work had shown that the visual responses of FEF neurons are typically not selective for the properties of stimuli [29]. However, before saccades were generated, a discrimination process occurred by which the activity of visually responsive cells in the FEF evolved to signal the location of the oddball target regardless of the visual feature (e.g. color or form) that distinguished it from the distractors.

During our studies on saccade target selection in the FEF of macaques, we found that the target selection signal that arises in the FEF had a number of interesting properties [19]. First, the initial response to an array of stimuli was consistently attenuated relative to the initial response to a single stimulus. This attenuation may have been attributable either to suppressive interactions among stimuli or to uncertainty about the location of the target. A similar finding has recently been made in the superior colliculus [30]. Second, target selection was achieved by a reduction of the activity evoked by the distractors (even though they were still in the receptive field), leaving only cells with the target in their receptive field with maximal activity. The current data do not indicate whether selection is accomplished through passive decay of distractor responses coupled with enhancement of the target response or through active suppression of the distractor responses [31•]. Third, the magnitude of the suppression of distractor-evoked activity varied with the proximity of the target to the receptive field, often being greater if the target flanked the receptive field. Behavioral studies have indicated that distractors close to a target are less likely to be selected than are more distant distractors under some [32], but not all [9], conditions; a center-surround selection mechanism has also been inferred in more complex cognitive decisions [33].

Effects of experience on target selection processes

Experts are more likely than novices to ignore conspicuous but non-informative elements of a visual image from their area of expertise [34]. This observation emphasizes the extent to which attention and gaze are under voluntary control. The degree of control and the rules by which it is expressed are being clarified by recent behavioral studies. Some studies indicate that salient stimuli in search displays attract attention automatically [35,36]; however, other work indicates that cognitive strategies can prevent the capture of attention by salient stimuli [3,37]. Other behavioral studies have demonstrated that search is

facilitated by implicit or explicit knowledge of the target properties [38] or location [39••].

We recently found that the neural selection process expressed in FEF can be modified profoundly by differences in cognitive strategy derived from experience [20••]. The findings from FEF described above were made in monkeys trained to search for an oddball stimulus regardless of the particular visual feature that defined it (e.g. both a red target among green distractors and a green target among red distractors). To perform the task, the monkeys had generalized a strategy of searching for the oddball stimulus. We found that when monkeys were given exclusive experience with one visual search array (e.g. only red among green), they adopted a strategy of ignoring stimuli that were distractors, even when those same stimuli became the oddball target in the complementary visual search array (e.g. green among red). In monkeys employing this behavioral strategy, we discovered that half of the visually responsive cells in FEF exhibited a suppressed response to the learned distractor as soon as the neurons responded. In other words, these neurons, which in earlier work had responded indiscriminantly to the target or distractors, now exhibited an apparent feature selectivity in their initial responses completely unlike what had been observed before in FEF. This finding may be related to earlier demonstrations of visual discrimination in prefrontal cortex based on instructions [40–42]. In summary, the physiological and anatomical data indicate that FEF may be regarded as a saliency map, a representation of the visual field in which the location of potential targets are registered, tagged by both feature properties and prior knowledge or expectations. What is the visual input to such a saliency map?


Visual analysis and selection

A visual image must be analyzed to provide information necessary to guide gaze. It is useful to distinguish different kinds of visual analyses. Beyond simply detecting the presence of visual stimuli, a higher level of analysis discriminates between alternative interpretations of a visual element at one retinal location. A still higher level of analysis compares visual elements across the visual field to select conspicuous or behaviorally relevant elements. Neural correlates of stimulus discrimination and stimulus selection have been investigated and are described below.

Stimulus discrimination

The relationship between visual neural responses and behavioral discrimination has been investigated extensively [43–45,46••]. Modeling efforts based on data from the middle temporal visual area (MT) indicate that perceptual decisions can be based on the weakly correlated activity of small populations of neurons, not all responding optimally, combined through a somewhat noisy process to signal one of the alternative directions of motion [47•]. A critical question is how do other centers in the brain ‘read out’ the activity of sensory cortex. It appears that the decision

Figure 1



Target selection and saccade production by FEF. **(a)** Comparison of activation of a visually responsive FEF neuron during visual search when the target of the search array was in the receptive field (solid line) and when only distractors of the search array were in the receptive field (dashed line). The locus of fixation is shown on a representative visual search display. Although the plots of activity were derived from one neuron, the visual search display shows receptive fields at two locations, one enclosing the target (solid semicircle) and the other enclosing a distractor (dashed semicircle), to illustrate the visual input to the neuron on the different types of trials. The initial visual response does not distinguish whether the target or distractors of the search array appeared in the receptive field. Over the next 100 ms, the activity evolves to signal whether or not the target was in the receptive field. Adapted from [21••]. **(b)** Stochastic growth of movement-related activity in FEF before saccades. Movement-related activity of a FEF neuron is plotted from trials that had three ranges of saccade latency. The plots of activity stop at the instant of saccade initiation, which is indicated by the arrows. Saccadic eye movements to the target are shown on the search display. Adapted from [73••].

rule depends on the context of the behavior. When a graded response is called for, then vector averaging appears to be used, but when categorical responses are required, a winner-take-all mechanism is employed [48,49••,50]. The balance of speed versus accuracy influences the decision rule: responses following short reaction times tend to reflect the outcome of an averaging decision, whereas responses following longer reaction times reflect a winner-take-all rule. This relationship is commonly observed for saccades made when multiple stimuli are presented [9].

Usually, a given stimulus arrangement supports a unique perceptual state. However, certain stimulus configurations that are perfectly discriminable support mutually exclusive perceptual states, such as the Necker cube or binocular rivalry. It has proven possible to relate the responses of neurons to ambiguous stimuli to the alternative perceptual

states reported by monkeys [51]. The correlation between neural activity and perceptual report improves as signals are conveyed further from the retina [52,53••]. These results indicate that when confronted with equally viable but competing interpretations of the image, the visual system does not tolerate ambiguity but adopts one of the alternative interpretations.

Stimulus selection

Visual elements can be selected on the basis of properties distinguishing them from surrounding elements or on the basis of prior instructions. Neural correlates of both kinds of visual selection have been described. Recent studies have demonstrated that a population of neurons in striate cortex is sensitive to local stimulus irregularities [54] and global texture gradients [55••]. In addition, a number of recent studies have investigated how the responses of cells in primary visual cortex (V1), extrastriate areas V2 and V4 [56••,57•], MT and MST [58], posterior parietal cortex (LIP and area 7a) [59•,60–62] and rostral inferior temporal cortex [63] are modulated according to the task instructions of what stimulus property or visual field location to attend to guide the behavioral response. Generally, responses to attended stimuli are greater than responses to unattended stimuli, accounting for intrinsic stimulus preferences. This modulatory effect appears to be more pronounced in areas further removed from V1 and is commonly greater if multiple stimuli are in a neuron's receptive field.

The mechanisms of the various forms of extraretinal modulation are not known. One informative avenue for elucidating the contributions of different mechanisms is examination of the time course of the modulatory effects both within the response of a single trial and across multiple trials as goals change. Unfortunately, many studies do not provide detailed information about the time course of the modulation for individual cells, although there are exceptions. For example, the long latency and spatial extent of the global texture effect observed in V1 [55••] suggests that it originates in feedback from extrastriate visual areas. Also, when stimuli were presented sequentially, the effects of attentional allocation were evident as soon as V4 neurons responded, but when multiple stimuli were flashed simultaneously, then attentional modulation was delayed slightly [56••]. Under both conditions, however, the attentional modulation of V4 neurons was multiphasic, with more pronounced modulation appearing 100–200 ms after stimulus presentation; such patterns are suggestive of different sources of modulation. Now, as we learn more about the time course of visual selection in different areas of the brain, it will be important to relate the different phases of selection to behavioral reaction times.

Time course of saccade target selection

The duration and variability of experimental reaction times or natural fixation periods exceed what can be


accounted for by simply adding transduction and transmission times from the sensory surface to the muscles [64]. Experimental psychology began with the hypothesis that reaction times are composed of stages of processing, and many behavioral studies have been conducted to investigate the flow of signals between these stages [65–67]. However, the conclusions of these behavioral studies are limited because they lack clear markers for the end of one stage and the beginning of another. Physiological measures can provide such markers. For instance, using event-related potentials (ERPs), investigators have deduced that the P300 component represents the conclusion of perceptual processing, and the lateralized readiness potential represents response preparation [68]. Unfortunately, the lack of spatial resolution and of knowledge of the neural generators of ERPs limit conclusions that can be drawn from these studies. To investigate stages of processing with the highest spatial and temporal resolution, single-neuron recording work can be guided by the insights of ERP studies inspired by psychological theories.

Having found that saccade target selection by FEF neurons occurred over time, we reasoned that a good marker for the conclusion of perceptual processing was the time at which FEF neurons registered whether the target or only distractors were in the response field [21••]. We related this time to variations in saccade latency and found that most visually responsive FEF neurons discriminated the target from distractors at a fairly constant interval after stimulus presentation (Figure 2). In other words, the time at which FEF neurons discriminated the target of the search array did not predict when gaze would shift to the target. In fact, FEF neurons discriminated the salient oddball stimulus from distractors even if monkeys never shifted gaze to the oddball target of a search array [22•]. Therefore, if the time of target discrimination by FEF neurons indexes the outcome and conclusion of perceptual processing, then it appears that although perceptual processing occupies a good portion of the reaction time, an additional delay and much of the variability of reaction time is unaccounted for. Thus, a postperceptual stage of response preparation must introduce the additional delay and variability observed in reaction times. This finding is consistent with conclusions from ERP studies [68] and substantiates the decoupling between perceptual processing and response preparation that has been observed in behavioral studies [11••,12]. This finding also suggests the possibility that the period after the target has been discriminated but before the saccade is produced may be the state in which attention has shifted to the location of the target.

From perception to action

Discrimination of sensory stimuli, while admittedly complicated and not fully understood, is still just the first step. If it is to guide action, one of the discriminated objects must be selected. But once an object is selected in the

Figure 2



Neural and behavioral chronometry. **(a)** Visual response latencies indicate when neurons in FEF became activated following presentation of the visual search array. The initial response did not discriminate the target from distractors. **(b)** The time of target discrimination indicates when the activity of FEF neurons reaches a state that is a statistically reliable signal of whether the target was in the response field. **(c)** Saccade initiation time measures when monkeys shifted gaze to fixate the selected target. During simple visual search, the time at which most FEF neurons discriminated the target did not predict when the eyes would move.

image, what is to be done with it? Approach or avoid? Look or not look? Grasp or not grasp? Because a given stimulus can afford different actions depending on experience and goals, it is necessary to distinguish the process that asks 'What is there?' from the process that asks 'What to do?'. The stage of processing in which the second question is answered is response selection and preparation. Several neurophysiological studies have investigated how particular stimuli become mapped onto arbitrary behaviors [69–71,72••]. The results of these studies uniformly demonstrate an evolution of neural activity during the reaction time from an early representation of the location of the sensory cue stimulus to a later representation of the direction of movement. Each study also identified neurons representing more complex contingencies of sensory and tonic activity on the stimulus–response mapping rule.

Once a behavioral response has been selected, the remaining decision to make is when to move. For gaze shifts, this is accomplished through the delicate balance of gaze-holding and gaze-shifting mechanisms in the oculomotor system. We have recently analyzed movement-related activity recorded in the FEF to evalu-

ate alternative models of reaction time [73••]. As illustrated in Figure 1b, we found that saccades were initiated when movement-related activity in the FEF reached a particular level that did not vary with saccade latency. Similar conclusions have been drawn from lateralized readiness potentials [74], from recordings in premotor and primary motor cortex [75], and in superior colliculus [76,77]. Thus, if the trigger mechanism for movement has a fixed threshold, then what can account for the stochastic variability of reaction time? In single-unit data from FEF [73••], the variability in reaction time appears to be accounted for by variation in the rate of growth of the premovement activity, which began at a fairly constant interval after target presentation, towards the threshold. In other words, trials with longer reaction times occurred when the rate of growth of the premovement activity was slower than on trials with shorter reaction times. The movement-related neural activity in FEF closely resembles the elements of models of reaction time [78,79].

Conclusions

Decisions transpire over time as multipotentiality gives way to determination. This review has described recent studies that have identified neural correlates of this transition in perceptual and postperceptual stages of processing. By emphasizing the chronometric approach, neuroscience and cognitive psychology can collaborate fruitfully to provide a fuller understanding of how the brain makes decisions. In all likelihood, the mechanisms revealed by studies of perceptual decisions may well apply to more portentous decisions.

Acknowledgements

We thank JH Kaas, SC Rao and K Thompson for their helpful comments on the manuscript. The work was supported by National Institutes of Health grants R01-EY08890, R01-MH55806, P30-EY08126. JD Schall is a Kennedy Center Investigator.

References and recommended reading

Papers of particular interest, published within the annual period of review, have been highlighted as:

- of special interest
 - of outstanding interest
1. Bierce A: *The Devil's Dictionary*. New York: Sagamore Press; 1957.
 2. Desimone R, Duncan J: **Neural mechanisms of selective visual attention**. *Annu Rev Neurosci* 1995, **18**:193-222.
 3. Egeth HE, Yantis S: **Visual attention: control, representation, and time course**. *Annu Rev Psychol* 1997, **48**:269-297.
 4. Duncan J, Humphreys G, Ward R: **Competitive brain activity in visual attention**. *Curr Opin Neurobiol* 1997, **7**:255-261.
 5. Schall JD: **Neural basis of saccade target selection**. *Rev Neurosci* 1995, **6**:63-85.
 6. Yarbus AL: *Eye Movements and Vision*. New York: Plenum Press; 1967.
 7. Viviani P: **Eye movements in visual search: cognitive, perceptual and motor control aspects**. In *Eye Movements and their Role in Visual and Cognitive Processes*. Edited by Kowler E. Amsterdam: Elsevier Science Publishers BV (Biomedical Division); 1990:353-393.
 8. Ballard DH, Hayhoe MM, Pelz JB: **Memory representations in natural tasks**. *J Cogn Neurosci* 1995, **7**:66-80.
 9. Findlay JM: **Saccade target selection during visual search**. *Vision Res* 1997, **37**:617-631.
 10. Zelinsky GJ, Sheinberg DL: **Eye movements during parallel-serial visual search**. *J Exp Psychol [Hum Percept Perform]* 1997, **23**:244-262.
 11. Hooge ITC, Erkelens CJ: **Control of fixation duration in a simple search task**. *Percept Psychophys* 1996, **58**:969-976.
This study demonstrates a dissociation between the timing of gaze shifts and visual processing. Although performance was characteristically idiosyncratic, the scanning eye movements of the subjects indicated that visual processing of the image centered on the fovea did not directly control saccade production.
 12. Kowler E, Anderson E, Doshier B, Blaser E: **The role of attention in the programming of saccades**. *Vision Res* 1995, **35**:1897-1916.
 13. Deubel H, Schneider WX: **Saccade target selection and object recognition: evidence for a common attentional mechanism**. *Vision Res* 1996, **36**:1827-1837.
A dual-task paradigm demonstrated that visual discrimination is best at the location of the intended target of a saccade, and directing attention to a discrimination target could not be done if a saccade was planned elsewhere. The data indicate that a common visual selection mechanism guides covert and overt orienting.
 14. Kustov AA, Robinson DL: **Shared neural control of attentional shifts and eye movements**. *Nature* 1996, **384**:74-77.
The study presents both microstimulation and single-neuron recording data from the superior colliculus. If monkeys attend, cued either exogenously or endogenously, to one location, then the eye movements evoked by microstimulation of the superior colliculus deviate toward the attended location. The effect had shorter latency and was more pronounced for exogenously than for endogenously cued attention shifts. The data also show that neurons in the superior colliculus are activated during both exogenously and endogenously cued attention shifts.
 15. Sheliga BM, Riggio L, Rizzolatti G: **Spatial attention and eye movements**. *Exp Brain Res* 1995, **105**:261-275.
 16. Nobre AC, Sebestyen GN, Gitelman DR, Mesulam MM, Frackowiak RSJ, Frith CD: **Functional localization of the system for visuospatial attention using positron emission tomography**. *Brain* 1997, **120**:515-533.
 17. Corbetta M: **Frontoparietal cortical networks for directing attention and the eye to visual locations: identical, independent or overlapping neural systems?** *Proc Natl Acad Sci USA* 1998, **95**:831-838.
 18. Meyer DE, Osman AM, Irwin DE, Yantis S: **Modern mental chronometry**. *Biol Psychol* 1988, **26**:3-67.
 19. Schall JD, Hanes DP, Thompson KG, King DJ: **Saccade target selection in frontal eye field of macaque. I. Visual and premovement activation**. *J Neurosci* 1995, **15**:6905-6918.
 20. Bichot NP, Schall JD, Thompson KG: **Visual feature selectivity in frontal eye fields induced by experience in mature macaques**. *Nature* 1996, **381**:697-699.
This paper reports an unexpected finding. The visual responses of many FEF neurons acquire feature selectivity if monkeys are trained to make saccades to the oddball target of only one of two complementary visual search arrays. Because the monkeys' task was the same in this condition as that of monkeys trained with both complementary search arrays, this initial visual selection cannot be attributable to motor enhancement. Also, because there could be no anticipation of target location on each trial, this initial visual selection cannot be attributable to spatial expectancy. Furthermore, the latency of the differential visual responses were too short to be explained by attentional selection. Prolonged repetition of one stimulus-response mapping rule may, therefore, lead to changes in functional connectivity in a feature map.
 21. Thompson KG, Hanes DP, Bichot NP, Schall JD: **Perceptual and motor processing stages identified in the activity of macaque frontal eye field neurons during visual search**. *J Neurophysiol* 1996, **76**:4040-4055.
Reaction time was partitioned into perceptual and motor components by relating the time at which FEF neurons discriminated the target from a distractor of a visual search array to saccade latency. Much of the variability of the saccade latencies was attributed to post-perceptual processes.
 22. Thompson KG, Bichot NP, Schall JD: **Dissociation of visual discrimination from saccade programming in macaque frontal eye field**. *J Neurophysiol* 1997, **77**:1046-1050.
Visually responsive neurons in FEF discriminated the target from the distractor of a visual search array even if monkeys never shifted gaze to the location occupied by the oddball target. These data are consistent with the possibility that FEF can play a role in covert attention shifts.

23. Burman DD, Segraves MA: **Primate frontal eye field activity during natural scanning eye movements.** *J Neurophysiol* 1994, **71**:1266-1271.
24. Schall JD, Morel A, King DJ, Bullier J: **Topography of visual cortical afferents to frontal eye field in macaque: functional convergence and segregation of processing streams.** *J Neurosci* 1995, **15**:4464-4487.
25. Segraves MA, Goldberg ME: **Functional properties of corticotectal neurons in the monkey's frontal eye fields.** *J Neurophysiol* 1987, **58**:1387-1419.
26. Segraves MA: **Activity of monkey frontal eye field neurons projecting to oculomotor regions of the pons.** *J Neurophysiol* 1992, **68**:1967-1985.
27. Dias EC, Kiesau M, Segraves MA: **Acute activation and inactivation of macaque frontal eye field with GABA-related drugs.** *J Neurophysiol* 1995, **74**:2744-2748.
28. Sommer MA, Tehovnik EJ: **Reversible inactivation of macaque frontal eye field.** *Exp Brain Res* 1997, **116**:229-249.
 This study and [27] demonstrate a profound incapacity of monkeys to generate eye movements following reversible inactivation of FEF. The results of these studies represent an important contrast to the numerous studies of the effects of lesions of FEF that have demonstrated that accurate saccades can be produced after a recovery period, probably through adaptive mechanisms of plasticity.
29. Mohler CW, Goldberg ME, Wurtz RH: **Visual receptive fields of frontal eye field neurons.** *Brain Res* 1973, **61**:385-389.
30. Basso MA, Wurtz RH: **Modulation of neural activity by target uncertainty.** *Nature* 1997, **389**:66-69.
31. Cepeda NJ, Cave KR, Bichot NP, Kim M-S: **Spatial selection via feature-driven inhibition of distractor locations.** *Percept Psychophys* 1998, in press.
 By comparing reaction times to probes at target, distractor and intervening locations, this study shows that attention in a search for a color target is primarily mediated by inhibition of the distractor locations.
32. Cave KR, Zimmerman JM: **Flexibility in spatial attention before and after practice.** *Psychol Sci* 1997, **8**:399-403.
33. Dagenbach D, Carr TH: **Inhibitory processes in perceptual recognition: evidence for a center-surround attentional mechanism.** In *Inhibitory Processes in Attention, Memory and Language*. Edited by Dagenbach D, Carr TH. New York: Academic Press; 1994:327-357.
34. Nodine CF, Kundel HL, Lauver SC, Toto LC: **Nature of expertise in searching mammograms for breast masses.** *Acad Radiol* 1996, **3**:1000-1006.
35. Theeuwes J: **Perceptual selectivity for color and form.** *Percept Psychophys* 1992, **51**:599-606.
36. Joseph JS, Optican LM: **Involuntary attentional shifts due to orientation differences.** *Percept Psychophys* 1996, **58**:651-665.
37. Bacon WF, Egeth HE: **Overriding stimulus-driven attentional capture.** *Percept Psychophys* 1994, **55**:485-496.
38. Maljkovic V, Nakayama K: **Priming of pop-out: I. Role of features.** *Mem Cogn* 1994, **22**:657-672.
39. Maljkovic V, Nakayama K: **Priming of pop-out: II. Role of position.** *Percept Psychophys* 1996, **58**:977-991.
 In a visual search task, subjects' attention was guided by one property (e.g. color) of visual elements but they reported a different property (e.g. shape) of the elements. This study shows that the history of the location of the target or the history of the feature guiding attention [38] influences the visual selection process: that is, reaction times are shorter if the target location or attention-guiding feature is repeated than if it switches on successive trials. This priming effect, which was not under conscious control and lasted over just 5-8 trials, seems to be the product of an implicit perceptual memory system. The visual system appears to treat the recent past as a reliable predictor of the immediate future.
40. Watanabe M: **Frontal units of the monkey coding the associative significance of visual and auditory stimuli.** *Exp Brain Res* 1992, **89**:233-247.
41. Sakagami M, Niki H: **Encoding of behavioral significance of visual stimuli by primate prefrontal neurons: relation to relevant task conditions.** *Exp Brain Res* 1994, **97**:423-436.
42. Boussaoud D, Wise SP: **Primate frontal cortex: neuronal activity following attentional versus intentional cues.** *Exp Brain Res* 1993, **95**:15-27.
43. Bradley A, Skottun BC, Ohzawa I, Sclar G, Freeman RD: **Visual orientation and spatial frequency discrimination: a comparison of single neurons and behavior.** *J Neurophysiol* 1987, **57**:755-772.
44. Vogels R, Orban GA: **How well do response changes of striate neurons signal differences in orientation: a study in the discriminating monkey.** *J Neurosci* 1990, **10**:3543-3558.
45. Britten KH, Shadlen MN, Newsome WT, Movshon JA: **The analysis of visual motion: a comparison of neuronal and psychophysical performance.** *J Neurosci* 1992, **12**:4745-4765.
46. Britten KH, Newsome WT, Shadlen MN, Celebrini S, Movshon JA: **A relationship between behavioral choice and the visual responses of neurons in macaque MT.** *Vis Neurosci* 1996, **13**:87-100.
 This is the latest in a series of investigations of the relation between the perceptual report of monkeys and the neural activity of neurons involved in generating the percept. The present paper describes the relation between neural activity and behavioral responses when the motion signal is absent or very close to threshold. In monkeys performing a two-alternative forced choice task, the variable responses of MT neurons to stochastic displays with no net motion showed a weak correlation with the behavioral report of the monkeys. Notably, though, the decision signal on trials with no net motion was not evident in the initial response of the population of neurons but arose some 40 ms after the activation had begun. A critical question is whether this delay arises from the time needed for the visual motion network to adopt one of the two possible states or whether it arises from the time needed for a top-down process to register the absence of clear evidence for either alternative and to mandate one of the possibilities.
47. Shadlen MN, Britten KH, Newsome WT, Movshon JA: **A computational analysis of the relationship between neuronal and behavioral responses to visual motion.** *J Neurosci* 1996, **16**:1486-1510.
 This paper sets a standard for developing theoretical relationships between perceptual reports and neural activity.
48. Ferrera VP, Lisberger SG: **Attention and target selection for smooth pursuit eye movements.** *J Neurosci* 1995, **14**:7472-7484.
49. Groh JM, Born RT, Newsome WT: **How is a sensory map read out? Effects of microstimulation in visual area MT on saccades and smooth pursuit eye movements.** *J Neurosci* 1997, **17**:4312-4330.
 Monkeys responded to slowly moving light spots by generating pursuit or saccadic eye movements. The accuracy of the eye movements depended on the estimation of the stimulus velocity. Normal performance was compared with performance when loci with known motion direction preferences in area MT were electrically microstimulated. The pattern of eye movement responses in most cases indicated that the velocity signal generated by the microstimulation was combined in a vectorial fashion with velocity signal derived from the visual stimulus. This evidence for a vector-average decision rule contrasts with a winner-take-all decision rule revealed by another study using microstimulation of area MT in monkeys performing an eight-alternative forced choice direction of motion judgement [50]. Evidently, the decision rule employed adapts to the opportunities and demands of the current situation.
50. Salzman CD, Newsome WT: **Neural mechanisms for forming a perceptual decision.** *Science* 1994, **264**:231-237.
51. Logothetis NK, Schall JD: **Neuronal correlates of subjective visual perception.** *Science* 1989, **245**:761-763.
52. Leopold DA, Logothetis NK: **Activity changes in early visual cortex reflect monkeys' percepts during binocular rivalry.** *Nature* 1996, **379**:549-553.
53. Sheinberg DL, Logothetis NK: **The role of temporal cortical areas in perceptual organization.** *Proc Natl Acad Sci USA* 1997, **94**:3408-3413.
 In previous studies, neural recordings from striate cortex and extrastriate visual areas such as V1, V2, V4 [52] and MT [51] during binocular rivalry found many neurons responding according to the retinal stimulus pattern and not the monkeys' perceptual reports. This study shows that the activity of neurons in higher-level temporal lobe visual areas in response to competing stimuli was highly correlated with the monkeys' perceptual report. The cortical representation at this stage of processing seems to correspond to the monkeys' perceptual state.
54. Knierim JJ, Van Essen DC: **Neuronal responses to static texture patterns in area V1 of the alert macaque monkey.** *J Neurophysiol* 1992, **67**:961-980.
55. Zipser K, Lamme VAF, Schiller PH: **Contextual modulation in primary visual cortex.** *J Neurosci* 1996, **16**:7376-7389.
 A long latency component of the responses of neurons in striate cortex to a texture patch covering their receptive field was enhanced if the texture surrounding the central patch was made perceptually distinct from the central

patch by differing in orientation, luminance, disparity, or color. This modulatory effect, which correlated with perceptual experience and extended up to 10 degrees in diameter, is an unexpected property of V1 neuron receptive fields.

56. Luck SJ, Chelazzi L, Hillyard SA, Desimone R: **Neural mechanisms of spatial selective attention in areas V1, V2, and V4 of macaque visual cortex.** *J Neurophysiol* 1997, **77**:24-42.

This paper describes a number of properties of the neural modulation associated with visual attention. The strength of modulation increased from V1 to V2 to V4. More consistent and stronger modulation was observed specifically when more than one stimulus appeared in the receptive field of the recorded neuron. When one location was to be attended across a block of trials, the spontaneous activity of V4 neurons representing the attended location increased before the stimuli appeared. This is one of the most extensive, systematic investigations of the neural correlates of visual attention that has been published.

57. Connor CE, Preddie DC, Gallant JL, Van Essen DC: **Spatial attention effects in macaque area V4.** *J Neurosci* 1997, **17**:3201-3214.

This paper reports the first clear evidence that focal visual attention influences the spatial distribution of visual processing.

58. Treue S, Maunsell JHR: **Attentional modulation of visual motion processing in cortical areas MT and MST.** *Nature* 1996, **382**:539-541.

59. Shadlen MN, Newsome WT: **Motion perception: seeing and deciding.** *Proc Natl Acad Sci USA* 1996, **93**:628-633.

Neural activity in area LIP was recorded in monkeys performing a two-alternative forced choice motion direction discrimination task. Monkeys responded with a saccade to one of two targets that corresponded to the perceived direction of motion of a random dot display centered on the fovea. In this study, unlike earlier studies [45,46**], one of the saccade targets rather than the motion display was placed in the neuron's receptive field. LIP neural activity reflected the discrimination process as it proceeded over time. An index of the sensory decision derived from the neural activity varied systematically with the strength of the motion signal, growing earlier and to a higher level the stronger the signal.

60. Robinson DL, Bowman EM, Kertzman C: **Covert orienting of attention in macaques. II. Contributions of parietal cortex.** *J Neurophysiol* 1995, **74**:698-712.

61. Steinmetz MA, Constantinidis C: **Neurophysiological evidence for a role of posterior parietal cortex in redirecting visual attention.** *Cereb Cortex* 1995, **5**:448-456.

62. Platt ML, Glimcher PW: **Responses of intraparietal neurons to saccadic targets and visual distractors.** *J Neurophysiol* 1997, **78**:1574-1589.

63. Chelazzi L, Miller EK, Duncan J, Desimone R: **A neural basis for visual search in inferior temporal cortex.** *Nature* 1993, **363**:345-347.

64. Carpenter RHS: **Oculomotor procrastination.** In *Eye Movements: Cognition and Visual Perception*. Edited by Fisher DF, Monty RA, Senders JW. Hillsdale, New Jersey: Lawrence Erlbaum; 1981:237-246.

65. Posner MI: *Chronometric Explorations of Mind*. Hillsdale, New Jersey: Lawrence Erlbaum Associates; 1978.

66. Luce RD: *Response Times: Their Role in Inferring Elementary Mental Organization*. Oxford: Oxford University Press; 1986.

67. Meyer DE, Osman AM, Irwin DE, Yantis S: **Modern mental chronometry.** *Biol Psychol* 1988, **26**:3-67.

68. Coles MGH, Smid HGOM, Scheffers MK, Otten LJ: **Mental chronometry and the study of human information processing.** In *Electrophysiology of Mind: Event-Related Brain Potentials and Cognition*. Edited by Rugg MD, Coles MGH. Oxford: Oxford University Press; 1995:86-131.

69. Di Pellegrino G, Wise SP: **Visuospatial versus visuomotor activity in the premotor and prefrontal cortex of a primate.** *J Neurosci* 1993, **13**:1227-1243.

70. Crammond DJ, Kalaska JF: **Modulation of preparatory neuronal activity in dorsal premotor cortex due to stimulus-response compatibility.** *J Neurophysiol* 1994, **71**:1281-1284.

71. Shen L, Alexander GE: **Neural correlates of a spatial sensory-to-motor transformation in primary motor cortex.** *J Neurophysiol* 1997, **77**:1171-1194.

72. Zhang J, Riehle A, Requin J, Kornblum S: **Dynamics of single neuron activity in monkey primary motor cortex related to sensorimotor transformation.** *J Neurosci* 1997, **17**:2227-2246.

Previous studies have described the sensory to motor evolution of neural activity observed when monkeys must produce an arbitrary movement in response to a particular stimulus. But it has proven difficult to describe the neural instantiation of the stimulus-response mapping rule. This paper applies a new analytical approach to identify whether neural activity recorded during stimulus-response compatible and incompatible trials can be attributed to sensory, motor or intermediate decision processes.

73. Hanes DP, Schall JD: **Neural control of voluntary movement initiation.** *Science* 1996, **274**:427-430.

Alternative theories of the origin of stochastic variability of reaction time were evaluated based on the movement-related activity of neurons in FEF. The evidence was entirely consistent with a variable-rate of accumulation and fixed-threshold architecture and contradicted a fixed-rate of accumulation and variable-threshold architecture. The distribution of reaction times generated by a fixed-threshold, variable-accumulator model of reaction time using parameters derived from single neurons corresponded in most cases to the distribution of reaction times actually produced by the monkey while that neuron was recorded.

74. Gratton G, Coles MGH, Sirevaag EJ, Eriksen CJ, Donchin E: **Pre- and poststimulus activation of response channels: a psychophysiological analysis.** *J Exp Psychol [Hum Percept Perform]* 1988, **14**:331-344.

75. Lecas J-C, Requin J, Anger C, Vitton N: **Changes in neuronal activity of the monkey precentral cortex during preparation for movement.** *J Neurophysiol* 1986, **56**:1680-1702.

76. Dorris MC, Paré M, Munoz DP: **Neuronal activity in monkey superior colliculus related to the initiation of saccadic eye movements.** *J Neurosci* 1997, **17**:8566-8579.

77. Stanford TR, Freedman EG, Sparks DL: **Site and parameters of microstimulation: evidence for independent effects on the properties of saccades evoked from the primate superior colliculus.** *J Neurophysiol* 1996, **76**:3360-3381.

78. Carpenter RHS, Williams MLL: **Neural computation of log likelihood in the control of saccadic eye movements.** *Nature* 1995, **377**:59-62.

79. Ratcliff R, Van Zandt T, McKoon G: **Connectionist and diffusion models of reaction time.** *Psychol Rev* 1998, in press.